

Alexander polynomials and Poincaré series of sets of ideals. *

A. Campillo F. Delgado[†] S.M. Gusein-Zade[‡]

In [2], [9], [4], [5], there were considered and, in some cases, computed Poincaré series of two sorts of multi-index filtrations on the ring of germs of functions on a complex (normal) surface singularity $(S, 0)$ (in particular on $(\mathbb{C}^2, 0)$). A filtration from the first class was defined by a curve (with several branches) on $(S, 0)$. For $(S, 0) = (\mathbb{C}^2, 0)$, the corresponding Poincaré series turned out to coincide with the Alexander polynomial (in several variables) of the corresponding (algebraic) link (when the number of branches is greater than 1; otherwise it is equal to the corresponding zeta function, i.e. to the Alexander polynomial in one variable divided by $1 - t$). Identifying all the variables, one gets the monodromy zeta function of the curve singularity. A filtration from the second class (so called divisorial one) was defined by a set of components of the exceptional divisor of a modification of the surface singularity $(S, 0)$. Here we define a notion of the Poincaré series of a set of ideals in the ring $\mathcal{O}_{(S, 0)}$ of germs of functions on $(S, 0)$, describe its connection with Poincaré series of filtrations and compute the Poincaré series of sets of ideals in some cases.

The discussed notion of the Poincaré series of a set of ideals was inspired by the notion of the zeta function of an ideal given in [15]. For our aim it is convenient to extend this notion to a finite set of ideals $\{I_1, \dots, I_r\}$, defining the “Alexander polynomial” of this set. This is a mixture of the notions introduced in [15] and [16].

*Math. Subject Class.: 14J17, 32S25. Keywords: ideals, surfaces, Poincaré series, zeta functions.

[†]First two authors were partially supported by the grant MTM2007-64704. Address: University of Valladolid, Dept. of Algebra, Geometry and Topology, 47011 Valladolid, Spain. E-mail: campillo@agt.uva.es, fdelgado@agt.uva.es

[‡]The research was partially supported by the grants RFBR-007-00593, INTAS-05-7805, and NWO-RFBR 047.011.2004.026. Address: Moscow State University, Faculty of Mathematics and Mechanics, Moscow, GSP-1, 119991, Russia. E-mail: sabir@mccme.ru

Let $(V, 0)$ be a germ of an analytic space with an isolated singular point at the origin and let $I_1, \dots, I_r$ be ideals in the ring $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$. Let $\pi : (X, D) \rightarrow (V, 0)$ be a resolution of the singularity of V and also of the set of the ideals $\{I_1, \dots, I_r\}$. This means that:

- 1) X is a complex analytic manifold;
- 2) π is a proper analytic map which is an isomorphism outside of the union of the zero loci of the ideals $I_1, \dots, I_r$;
- 3) the exceptional divisor $D = \pi^{-1}(0)$ is a normal crossing divisor on X ;
- 4) for $i = 1, \dots, r$, the lifting $I_i^* = \pi^* I_i$ of the ideal I_i to the space X of the modification is locally principal (and therefore π is a principalization of the ideal I_i);
- 5) the union of the zero loci of the liftings of the ideals $I_1, \dots, I_r$ to the space X of the modification is a normal crossing divisor on X .

For $\underline{k} = (k_1, \dots, k_r) \in \mathbb{Z}_{\geq 0}^r$, let $S_{\underline{k}}$ be the set of points $x \in D$ such that, in some local coordinates $z_1, \dots, z_n$ on X centred at the point x , the zero locus of the ideal I_{ix}^* , $i = 1, \dots, r$, is the hypersurface $\{z_1 = 0\}$ with multiplicity k_i , i.e. $I_{ix}^* = \langle z_1^{k_i} \rangle$.

Definition: *The Alexander polynomial of the set of ideals $\{I_i\}$ is the rational function (or a power series) in the variables $t_1, \dots, t_r$ given by the A'Campo type ([1]) formula:*

$$\Delta_{\{I_i\}}(t_1, \dots, t_r) = \prod_{\underline{k} \in \mathbb{Z}_{\geq 0}^r \setminus \{0\}} (1 - \underline{t}^{\underline{k}})^{-\chi(S_{\underline{k}})},$$

where $\chi(\cdot)$ is the Euler characteristic and $\underline{t}^{\underline{k}} := t_1^{k_1} \cdot \dots \cdot t_r^{k_r}$.

Remarks. 1. For $r = 1$, this gives the notion of the zeta function $\zeta_I(t)$ of an ideal I : [15]. In [15, Theorem 4.2] one forgot to write that π should be an isomorphism outside of the zero locus of the ideal I . This is the reason why we formulate the notion for germs $(V, 0)$ with isolated singularities. Another option is to demand that the singular locus of V is contained in the zero locus of the ideal I .

2. The Alexander polynomial defined this way is not, in general, a polynomial. It is really a polynomial in the case $V = \mathbb{C}^2$ and $I_i = \langle f_i \rangle$, where $f_i = 0$ are equations of the irreducible components C_i of a (reduced) plane curve singularity $(C, 0) \subset (\mathbb{C}^2, 0)$: $C = \bigcup_{i=1}^r C_i$, $r > 1$. In this case $\Delta_{\{I_i\}}(t_1, \dots, t_r)$

coincides with the classical Alexander polynomial in several variables of the algebraic link $C \cap S_\varepsilon^3 \subset S_\varepsilon^3$ (S_ε^3 is the sphere of small radius ε centred at the origin in $\mathbb{C}^2$): see [10]. We prefer to keep the name in the general case.

Let $(V, 0)$ be a germ of an analytic space and let $\mathbb{P}\mathcal{O}_{V,0}$ be the projectivization of the ring $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$. The following concept was described, e.g., in [3].

Let $J_{V,0}^k = \mathcal{O}_{V,0}/\mathfrak{m}^{k+1}$ be the space of k -jets of functions on $(V; 0)$ ($\mathfrak{m}$ is the maximal ideal in the ring $\mathcal{O}_{V,0}$), let $\mathbb{P}J_{V,0}^k$ be its projectivization, and let $\mathbb{P}^*J_{V,0}^k := \mathbb{P}J_{V,0}^k \cup \{*\}$. Let $\pi_k : \mathbb{P}\mathcal{O}_{V,0} \rightarrow \mathbb{P}^*J_{V,0}^k$ be the natural map. A subset $A \subset \mathbb{P}\mathcal{O}_{V,0}$ is said to be cylindric if $A = \pi_k^{-1}(B)$ for a constructible subset $B \subset \mathbb{P}J_{V,0}^k \subset \mathbb{P}^*J_{V,0}^k$. The Euler characteristic $\chi(A)$ of a cylindric set $A \subset \mathbb{P}\mathcal{O}_{V,0}$, $A = \pi_k^{-1}(B)$, is defined as the Euler characteristic of the set B . A function $\psi : \mathbb{P}\mathcal{O}_{V,0} \rightarrow G$ with values in an Abelian group G is called cylindric if, for each $a \in G$, $a \neq 0$, the set $\psi^{-1}(a)$ is cylindric. The integral $\int_{\mathbb{P}\mathcal{O}_{V,0}} \psi d\chi$ of a cylindric function ψ over the space $\mathbb{P}\mathcal{O}_{V,0}$ with respect to the Euler characteristic is the sum $\sum_{a \in G \setminus \{0\}} \chi(\psi^{-1}(a))a$ (if this sum makes sense in the group G).

Let

$$\mathcal{O}_{V,0} = J(0) \supset J(1) \supset J(2) \supset \dots \quad (1)$$

be a filtration of the ring $\mathcal{O}_{V,0}$ by $\mathbb{C}^*$ -invariant subsets $J(k)$ (not necessarily vector subspaces) with cylindric projectivizations $\mathbb{P}J(k) \subset \mathbb{P}\mathcal{O}_{V,0}$.

Definition: The Poincaré series of the filtration (1) is the series

$$P(t) = \sum_{k=0}^{\infty} \chi(\mathbb{P}J(k) \setminus \mathbb{P}J(k+1)) \cdot t^k \in \mathbb{Z}[[t]].$$

This definition can be rewritten as $P(t) = \int_{\mathbb{P}\mathcal{O}_{V,0}} t^{v(g)} d\chi$ where t^∞ is assumed to be equal to zero.

Example. If all the elements $J(k)$ of the filtration are vector subspaces of $\mathcal{O}_{V,0}$, then

$$P(t) = \sum_{k=0}^{\infty} \dim(J(k) \setminus J(k+1)) \cdot t^k.$$

A filtration like (1) may be defined by the corresponding ($\mathbb{C}^*$ -invariant) tautological function $v : \mathcal{O}_{V,0} \rightarrow \mathbb{Z}_{\geq 0} \cup \{+\infty\}$: $v(g) = \sup\{k : g \in J(k)\}$, $J(k) = \{g \in \mathcal{O}_{V,0} : v(g) \geq k\}$. The function v is cylindric in the sense that, for each $k \in \mathbb{Z}_{\geq 0}$, the set $\mathbb{P}J(k)$ is cylindric.

Let $\{v_1, \dots, v_r\}$ be a set of $\mathbb{C}^*$ -invariant cylindric functions on $\mathcal{O}_{V,0}$ with values in $\mathbb{Z}_{\geq 0} \cup \{+\infty\}$. This set defines a multi-index filtration on $\mathcal{O}_{V,0}$: for $\underline{v} = (v_1, \dots, v_r) \in \mathbb{Z}_{\geq 0}^r$,

$$J(\underline{v}) = \{g \in \mathcal{O}_{V,0} : \underline{v}(g) \geq \underline{v}\} \quad (2)$$

($\underline{v}(g) := (v_1(g), \dots, v_r(g))$, $\underline{v}' \geq \underline{v}''$ iff $v'_i \geq v''_i$ for all i). The sets $J(\underline{v})$ are not, in general, vector subspaces of $\mathcal{O}_{V,0}$. The sets $\mathbb{P}J(\underline{v})$ are cylindric.

Definition: The Poincaré series of the filtration (2) is the series

$$\begin{aligned} P(t_1, \dots, t_r) &= \int_{\mathbb{P}\mathcal{O}_{V,0}} \underline{t}^{\underline{v}(g)} d\chi \\ &= \sum_{\underline{v} \in \mathbb{Z}_{\geq 0}^r} \chi(\mathbb{P}J(\underline{v}) \setminus \bigcup_{i=1}^r \mathbb{P}J(\underline{v} + \underline{1}_i)) \cdot \underline{t}^{\underline{v}} \in \mathbb{Z}[[t_1, \dots, t_r]]. \end{aligned} \quad (3)$$

where $\underline{1}_i := (0, \dots, 0, 1, 0, \dots, 0)$ (1 is at the i th place), $\underline{t}^{\underline{v}} := t_1^{v_1} \cdot \dots \cdot t_r^{v_r}$.

Example. If all v_i , $i = 1, \dots, r$, are order functions, that is $v_i(g_1 + g_2) \geq \min\{v_i(g_1), v_i(g_2)\}$, all the elements $J(\underline{v})$ of the filtration are vector subspaces of $\mathcal{O}_{V,0}$, and the definition (3) of the Poincaré series becomes equivalent to the definition in [2, 3] given in terms of dimensions.

Now let $(S, 0)$ be a germ of a normal surface singularity. For a function germ $g \in \mathcal{O}_{S,0}$ and a divisor γ on $(S, 0)$, there is defined the intersection number $(\gamma \circ g) \in \mathbb{Z} \cup \{\infty\}$. It can be defined, e.g., in the following way. Let $\gamma = \sum_i k_i \gamma_i$ where γ_i are divisors represented by irreducible curves $(C_i, 0)$ on $(S, 0)$. Then $(\gamma \circ g) = \sum_i k_i (\gamma_i \circ g)$ where $(\gamma_i \circ g)$ is the order of zero of the function g on the curve $(C_i, 0)$ in a uniformization parameter. (If $g|_{C_i} \equiv 0$, $(\gamma_i \circ g)$ is assumed to be equal to $+\infty$.) If γ is a Cartier divisor, $\gamma = \{f = 0\}$, $f \in \mathcal{O}_{S,0}$, we shall write $(f \circ g)$ instead of $(\gamma \circ g)$.

Let I be an ideal in the ring $\mathcal{O}_{S,0}$ and let $v_I(g) := \min\{(g \circ f) : f \in I\}$. One can see that the function $v_I : \mathcal{O}_{S,0} \rightarrow \mathbb{Z}_{\geq 0} \cup \{\infty\}$ is $\mathbb{C}^*$ -invariant and cylindric.

Definition: The filtration corresponding to the ideal I is the filtration

$$\mathcal{O}_{S,0} = J_I(0) \supset J_I(1) \supset J_I(2) \supset \dots$$

defined by the function v_I :

$$J_I(k) = \{g \in \mathcal{O}_{S,0} : v_I(g) \geq k\}.$$

Now let $\{I_1, \dots, I_r\}$ be a set of ideals in the ring $\mathcal{O}_{S,0}$.

Definition: The multi-index filtration corresponding to the set of ideals $\{I_1, \dots, I_r\}$ is the filtration of $\mathcal{O}_{S,0}$, the elements of which are

$$J_{\{I_i\}}(\underline{v}) = \bigcap_{i=1}^r J_{I_i}(v_i)$$

($\underline{v} = (v_1, \dots, v_r) \in \mathbb{Z}_{\geq 0}^r$).

This filtration is defined by the set $\{v_{I_i}\}$ of the corresponding functions. The spaces $J_{\{I_i\}}(\underline{v})$ are not, in general, vector subspaces of $\mathcal{O}_{S,0}$. One can see that the subsets $J_{\{I_i\}}(\underline{v})$ are, so called, "linear subspaces", i.e. belong to the algebra generated by vector subspaces.

Examples. 1. Let $(C, 0) \subset (S, 0)$ be a germ of a (reduced) curve on the surface S , let $C = \bigcup_{i=1}^r C_i$ be the decomposition of the curve C into irreducible components, let I_{C_i} be the ideal of the curve C_i . One has the multi-index filtration corresponding to the set of ideals $\{I_{C_1}, \dots, I_{C_r}\}$. This is a filtration of the ring $\mathcal{O}_{S,0}$ by ideals. For $S = \mathbb{C}^2$ or if all the components C_i of the curve C are Cartier divisors on $(S, 0)$ (this takes place, e.g., for any curve on the rational double point of type E_8), this is the filtration corresponding to the curve C considered in [2] and [5]. Otherwise this is not, generally speaking, the case. For $S = \mathbb{C}^2$, the Poincaré series of this filtration coincides with the Alexander polynomial (in r variables) of the curve $(C, 0)$, i.e. of the corresponding link $C \cap S_\varepsilon^3 \subset S_\varepsilon^3$: [2].

2. Let $\pi : (X, D) \rightarrow (S, 0)$ be a proper modification of the space $(S, 0)$ which is an isomorphism outside of the origin, with X smooth and $D = \pi^{-1}(0)$ a normal crossing divisor on X . Let $D = \bigcup_{\sigma \in \Gamma} E_\sigma$ be the representation of the exceptional divisor D as the union of its irreducible components. For $\sigma \in \Gamma$, i.e. for a component E_σ of the exceptional divisor D , let $\tilde{L}$ be a germ of a smooth irreducible curve on X intersecting E_σ transversally at a smooth point (i.e. not at an intersection point with other components of the exceptional divisor D), let $L = \pi(\tilde{L})$ be the corresponding curve on $(S, 0)$, let $I_L \subset \mathcal{O}_{S,0}$ be the ideal of the curve $(L, 0)$, and let I_{E_σ} be the ideal generated by all the ideals I_L of the described type. For r chosen components $E_1, \dots, E_r$ of the exceptional divisor D , i.e. for $\{1, \dots, r\} \subset \Gamma$, this way one gets a multi-index filtration (by ideals) corresponding to a set $\{E_1, \dots, E_r\}$ of components of the exceptional divisor D . Again, if $S = \mathbb{C}^2$ or if all curves L described above are Cartier divisors on $(S, 0)$, this filtration coincides with the divisorial filtration studied in [9] and [4]. Otherwise this is not the case. For $S = \mathbb{C}^2$, the Poincaré

series of this filtration coincides with the Alexander polynomial of the set of ideals $\{I_{E_1}, \dots, I_{E_r}\}$: [9].

3. One can consider a set of valuations $v_1, \dots, v_{r'}, w_1, \dots, w_{r''}$ corresponding to components $E_1, \dots, E_{r'}$ of a modification $\pi : (X, D) \rightarrow (S, 0)$ and to irreducible curve germs $(C_i, 0) \subset (S, 0)$, $i = 1, \dots, r''$, respectively. If $S = \mathbb{C}^2$, and the modification π is an embedded resolution of the curve $C = \bigcup_{i=1}^{r''} C_i$, one has the following formula for the Poincaré series of the corresponding multi-index filtration (by ideals). Let $m_{\sigma\delta}$ be the minus inverse matrix of the intersection matrix $(E_\sigma \circ E_\delta)$ of the components E_σ on X , let $\alpha_i \in \Gamma$, $i = 1, \dots, r''$, be the number of the component E_{α_i} of the exceptional divisor D intersecting the strict transform of the (irreducible) curve C_i , and let $\underline{m}'_\sigma := (m_{\sigma 1}, \dots, m_{\sigma r'})$, $\underline{m}''_\sigma := (m_{\sigma \alpha_1}, \dots, m_{\sigma \alpha_{r''}})$. Let $\overset{\circ}{E}_\sigma$ be the smooth part of the component E_σ in the union of preimages (total transforms) of the curves C_i , i.e. E_σ minus intersection points with other components of the exceptional divisor D and with the strict transforms of the curves C_i , $i = 1, \dots, r''$. Just as in [3] and [9] (see also the proof of Theorem 1), one can show that

$$P(t_1, \dots, t_{r'}, T_1, \dots, T_{r''}) = \prod_{\sigma \in \Gamma} (1 - \underline{t}^{\underline{m}'_\sigma} \underline{T}^{\underline{m}''_\sigma})^{-\chi(\overset{\circ}{E}_\sigma)}$$

(here $\underline{t} = (t_1, \dots, t_{r'})$ and $\underline{T} = (T_1, \dots, T_{r''})$ are the variables corresponding to the valuations $v_1, \dots, v_{r'}$ and $w_1, \dots, w_{r''}$ respectively).

Let $\{I_1, \dots, I_r\}$ be a set of ideals in $\mathcal{O}_{\mathbb{C}^2, 0}$. Let $\pi : (X, D) \rightarrow (\mathbb{C}^2, 0)$ be a resolution of the set of ideals $\{I_i\}$. Let $D = \bigcup_{\sigma \in \Gamma} E_\sigma$, where E_σ are irreducible components of the exceptional divisor D (each E_σ is isomorphic to the complex projective line $\mathbb{CP}^1$). For $\sigma \in \Gamma$, i.e. for a component E_σ of the exceptional divisor D , let $\overset{\circ}{E}_\sigma$ be the “smooth part” of E_σ in the union of zero loci of the ideals $\{I_i^*\}$, i.e. E_σ minus intersection points with all other components of the union. Let $k_{\sigma i}$ be the multiplicity of the component E_σ in the zero divisor of the ideal I_i and let $\underline{k}_\sigma := (k_{\sigma 1}, \dots, k_{\sigma r}) \in \mathbb{Z}_{\geq 0}^r$.

Theorem 1. *One has*

$$P_{\{I_i\}}(t_1, \dots, t_r) = \prod_{\sigma \in \Gamma} (1 - \underline{t}^{\underline{k}_\sigma})^{-\chi(\overset{\circ}{E}_\sigma)}. \quad (4)$$

This statement generalizes those from [2] and [9] for the ideals described in Examples 1 and 2 (for $S = \mathbb{C}^2$). One can see that the right-hand side of the equation (4) is equal to the Alexander polynomial $\Delta_{\{I_i\}}(t_1, \dots, t_r)$ of the set of ideals $\{I_i\}$ and thus in this case the Poincaré series coincides with the Alexander polynomial.

Proof. The proof essentially repeats the arguments from [3], [9]. One uses the representation of the Poincaré series as an integral with respect to the Euler characteristic. There is a map from the projectivization $\mathbb{P}\mathcal{O}_{\mathbb{C}^2,0}$ of the ring $\mathcal{O}_{\mathbb{C}^2,0}$ onto the space of effective divisors on $\mathring{D} = \bigcup_{\sigma \in \Gamma} \mathring{E}_\sigma$: to a function germ f on $(\mathbb{C}^2, 0)$ one associates the intersection of the strict transform of the curve $\{f = 0\}$ with the exceptional divisor D . (For any chosen N this map is defined for all function germs f with $v_{I_i}(f) \leq N$, $i = 1, \dots, r$, if one makes sufficiently many additional blowing-ups at intersection points of the components of the union of the zero loci of the ideals I_i^* on X . For all additional components E_σ of the exceptional divisor D , their smooth parts $\mathring{E}_\sigma$ are isomorphic to the complex projective line $\mathbb{CP}^1$ without two points. Therefore their Euler characteristics are equal to zero and they do not contribute to the right-hand side of the equation (4).) Proposition 2 from [3] implies that the preimage of a point with respect to this map is a complex affine space and thus has the Euler characteristic equal to 1. The Fubini formula implies that the Poincaré series $P_{\{I_i\}}(t_1, \dots, t_r)$ is equal to the integral with respect to the Euler characteristic of the monomial $\underline{t}^{\underline{v}}$ over the space of effective divisors on $\mathring{D}$. Here $\underline{v}$ is an additive function on the space of effective divisors on $\mathring{D}$ (with values in $\mathbb{Z}_{\geq 0}^r$) equal to $\underline{k}_\sigma$ for a point from the component $\mathring{E}_\sigma$.

The space of effective divisors on $\mathring{D}$ is the direct product of the spaces of effective divisors on the components $\mathring{E}_\sigma$, $\sigma \in \Gamma$. Each of the latter ones is the disjoint union of the symmetric powers $S^\ell \mathring{E}_\sigma$ of the component $\mathring{E}_\sigma$. Therefore

$$P_{\{I_i\}}(t_1, \dots, t_r) = \prod_{\sigma \in \Gamma} \left(\sum_{\ell=0}^{\infty} \chi(S^\ell \mathring{E}_\sigma) \cdot \underline{t}^{\ell \underline{k}_\sigma} \right).$$

Now equation (4) follows from the formula

$$\sum_{\ell=0}^{\infty} \chi(S^\ell X) t^\ell = (1 - t)^{-\chi(X)}.$$

□

Example 4. By the definition, for ideals $I_1, \dots, I_r \subset \mathcal{O}_{\mathbb{C}^2,0}$, their Alexander polynomial coincides with the Alexander polynomial of their integral closures $\bar{I}_1, \dots, \bar{I}_r$. An integrally closed ideal $I = \bar{I}$ has a representation of the form $I = \prod_{\sigma} I_{E_\sigma}^{n_\sigma} \prod_{j=1}^{r''} I_{C_j}^{m_j}$ where I_{E_σ} is the ideal corresponding to a divisor E_σ of a

resolution of the ideal I and I_{C_j} is the ideal of an irreducible curve singularity $(C_j, 0) \subset (\mathbb{C}^2, 0)$, $m_j > 0$ (see [17]). Let $s = \#\Gamma$ be the number of components of the exceptional divisor D of the resolution. One has $n_\sigma = \sum_{\delta \in \Gamma} m_{\sigma\delta} k_\delta$, where k_δ is the multiplicity of the ideal I on the component E_δ of the exceptional divisor D , $(m_{\sigma\delta})$ is the minus inverse matrix of the intersection matrix $(E_\sigma \circ E_\delta)$ of the components E_σ on X . Therefore one has the following equation for the order function $v_I : \mathcal{O}_{\mathbb{C}^2, 0} \setminus \{0\} \rightarrow \mathbb{Z}_{\geq 0}^r \cup \{\infty\}$:

$$v_I = \sum_{\sigma} n_{\sigma} v_{\sigma} + \sum_{j=1}^{r''} m_j w_j = \sum_{\sigma, \delta} m_{\sigma\delta} k_{\delta} v_{\sigma} + \sum_{j=1}^{r''} m_j w_j$$

where v_{σ} is the order function corresponding to the component E_{σ} (Example 2), w_j is the order function corresponding to the irreducible curve $(C_j, 0)$ (Example 1). One has the following equation

$$P_I(t) = P(t^{n_1}, \dots, t^{n_s}, t^{m_1}, \dots, t^{m_{r''}})$$

where $P(t_1, \dots, t_s, T_1, \dots, T_{r''})$ is the Poincaré series of the valuations $(v_1, \dots, v_s, w_1, \dots, w_{r''})$ corresponding to the components E_{σ} of the exceptional divisor and to the curves $(C_j, 0)$ (Example 3). Moreover, one has the following statement.

Proposition 1. *Let $(I_1, \dots, I_r)$ be a set of ideals in $\mathcal{O}_{\mathbb{C}^2, 0}$, and let $I_i = \prod_{\sigma} I_{\sigma}^{n_{\sigma}^i} \prod_{j=1}^{r''} I_{C_j}^{m_j^i}$, $i = 1, \dots, r$, where, for any $j = 1, \dots, r''$, there exists i with $m_j^i > 0$. Then one has*

$$P_{\{I_i\}}(t_1, \dots, t_r) = P\left(\prod_{i=1}^r t_i^{n_1^i}, \dots, \prod_{i=1}^r t_i^{n_s^i}, \prod_{i=1}^r t_i^{m_1^i}, \dots, \prod_{i=1}^r t_i^{m_{r''}^i}\right).$$

In a similar way one can prove versions of the main statements from [4] and [7] for ideals in the ring of functions on a rational surface singularity or on their universal abelian covers. Let $(S, 0)$ be a rational surface singularity and let $\pi : (X, D) \rightarrow (S, 0)$ be its resolution. The link $S \cap S_{\varepsilon}^3$ of the singularity $(S, 0)$ is a rational homology sphere and its first homology group $H = H_1(S \setminus \{0\})$ is isomorphic to the cokernel $\mathbb{Z}^{\Gamma} / \text{Im } j$ of the map $j : \mathbb{Z}^{\Gamma} \rightarrow \mathbb{Z}^{\Gamma}$ defined by the intersection matrix $(E_{\sigma} \circ E_{\sigma'})$ (the order of the group H is equal to the determinant d of minus the intersection matrix $-(E_{\sigma} \circ E_{\sigma'})$). For $\sigma \in \Gamma$, let h_{σ} be the element of the group H represented by the loop in the manifold $X \setminus D \simeq S \setminus \{0\}$ going around the component E_{σ} in the positive direction. The group H is generated by the elements h_{σ} for all $\sigma \in \Gamma$.

Let $p : (\tilde{S}, 0) \rightarrow (S; 0)$ be the universal abelian cover of the surface singularity S (see e.g. [13], [14], [12]). The group H acts on $(\tilde{S}, 0)$ and the restriction $p|_{\tilde{S} \setminus \{0\}}$ of the map p to the complement of the origin is a (usual, nonramified) covering $\tilde{S} \setminus \{0\} \rightarrow S \setminus \{0\}$ with the structure group H . One can lift the map p to a (ramified) covering $p' : (\tilde{X}, \tilde{D}) \rightarrow (X, D)$ where $\tilde{X}$ is a normal surface (generally speaking, not smooth) and $\tilde{X} \setminus \tilde{D} \simeq \tilde{S} \setminus \{0\}$:

$$\begin{array}{ccc} (\tilde{X}, \tilde{\mathcal{D}}) & \xrightarrow{\tilde{\pi}} & (\tilde{\mathcal{S}}, 0) \\ \downarrow p' & & \downarrow p \\ (X, \mathcal{D}) & \xrightarrow{\pi} & (\mathcal{S}, 0) \end{array}$$

(one can define $\tilde{X}$ as the normalization of the fibre product $X \times_{\mathcal{S}} \tilde{\mathcal{S}}$ of the varieties X and $\tilde{\mathcal{S}}$ over $\mathcal{S}$).

Let $R(H)$ be the ring of (virtual) representations of the group H . For $\sigma \in \Gamma$, i.e. for a component E_σ of the exceptional divisor D , let α_σ be the one-dimensional representation $H \rightarrow \mathbb{C}^* = \mathbf{GL}(1, \mathbb{C})$ of the group H defined by $\alpha_\sigma(h_\delta) = \exp(-2\pi\sqrt{-1}m_{\sigma\delta})$ (here the minus sign reflects the fact that the action of an element $h \in H$ on the ring $\mathcal{O}_{\tilde{S},0}$ is defined by $(h \cdot f)(x) = f(h^{-1}(x))$). Let $\{I_1, \dots, I_r\}$ be a set of ideals in $\mathcal{O}_{S,0}$ and let $\tilde{I}_i = p^*I_i$ be the liftings of the ideals I_i to the universal abelian cover $\tilde{S}$. The corresponding multi-index filtration on the ring $\mathcal{O}_{\tilde{S},0}$ of germs of functions on the abelian cover $\tilde{S}$ is an H -invariant one. A notion of the equivariant Poincaré series of such multi-index filtration was defined in [6]. Similar to [7] one obtains the following result.

Theorem 2. *If the resolution $\pi : (X, D) \rightarrow (S, 0)$ is a resolution of the set of ideals $\{I_1, \dots, I_r\}$, one has*

$$P^H(t_1, \dots, t_r) = \prod_{\sigma \in \Gamma} (1 - \alpha_\sigma t^{\underline{d}_{\sigma}})^{-\chi(\overset{\circ}{E}_\sigma)}. \quad (5)$$

The sum of monomials of this series with the trivial representation in the coefficients (i.e. those for which $\sum_{\sigma} m_{\delta\sigma} v_\sigma$ is an integer for any $\delta \in \Gamma$) with the change of variables $t_i^d \rightarrow t_i$ is the Poincaré series of the filtration on $\mathcal{O}_{S,0}$ corresponding to the set of ideals $\{I_i\}$ (cf. [4], [5]).

Remark. Integrally closed ideals in the ring $\mathcal{O}_{S,0}$ of germs of functions on a rational surface singularity $(S, 0)$ have a description somewhat similar to that in Example 4 for $S = \mathbb{C}^2$. Let $I = \bar{I}$ be an integrally closed ideal in $\mathcal{O}_{S,0}$ and let $\pi : (X, 0) \rightarrow (S, 0)$ be a resolution of it. For $\sigma \in \Gamma$, i.e. for a component E_σ of the exceptional divisor D , let $\tilde{L}$ be a germ of a smooth irreducible curve

on X intersecting E_σ transversally at a smooth point, i.e. at a point of $\overset{\circ}{E}_\sigma$, and let $L = \pi(\tilde{L})$. There exists the minimal natural number d_σ such that $d_\sigma L$ is a Cartier divisor on $(S, 0)$: $d_\sigma L = (g_L)$ for $g_L \in \mathcal{O}_{S,0}$. (The number d_σ is the minimal natural number such that $d_\sigma m_{\sigma\delta}$ are integers for all $\delta \in \Gamma$ and is equal to the order of the element h_σ in the group $H = H_1(S \setminus \{0\})$.) Let $I'_\sigma \subset \mathcal{O}_{S,0}$ be the ideal generated by all germs g_L of the described type.

The integrally closed ideal I has a unique representation of the form $I = \prod_{\sigma} I_{\sigma}^{r_{\sigma}}$ for non negative rational numbers r_{σ} such that $\sum_{\sigma} (E_{\sigma} \circ E_{\delta}) r_{\sigma}$ are integers for all $\delta \in \Gamma$ (see [11]). (For integrally closed ideals I_1 and I_2 in the ring $\mathcal{O}_{S,0}$, one writes that $I_1 = I_2^{1/q}$ if and only if $I_1^q = I_2$. If such an ideal I_1 exists, it is defined in a unique way.) As it was mentioned, the divisorial filtration on the ring $\mathcal{O}_{S,0}$ corresponding to the component E_σ is not, generally speaking, the filtration corresponding to an ideal. However, for the corresponding functions one has $v_{I'_\sigma} = d_\sigma v_{E_\sigma}$. Therefore the Poincaré series of the filtration corresponding to the ideal I'_σ is obtained from the one of the divisorial filtration corresponding to the component E_σ by substituting the variable t by t^{d_σ} . This explains a relation between the Poincaré series of the filtration in the ring $\mathcal{O}_{S,0}$ corresponding to a set of ideals and that of the divisorial filtration like in Example 4.

References

- [1] A'Campo N. La fonction zêta d'une monodromie. *Comment. Math. Helv.*, v.50, 233–248 (1975).
- [2] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity via the ring of functions on it. *Duke Math. J.*, v.117, no.1, 125–156 (2003).
- [3] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity and integrals with respect to the Euler characteristic. *Internat. J. Math.*, v.14, no.1, 47–54 (2003).
- [4] Campillo A., Delgado F., Gusein-Zade S.M. Poincaré series of a rational surface singularity. *Invent. math.*, v.155, 41–53 (2004).
- [5] Campillo A., Delgado F., Gusein-Zade S.M. Poincaré series of curves on rational surface singularities. *Commen. Math. Helv.*, v.80, no.1, 95–102 (2005).

- [6] Campillo A., Delgado F., Gusein-Zade S.M. On Poincaré series of filtrations on equivariant functions of two variables. *Moscow Math. J.*, v.7, no.2, 243–255 (2007).
- [7] Campillo A., Delgado F., Gusein-Zade S.M. Universal abelian covers of rational surface singularities and multi-index filtrations. *Funct. Anal. and Appl.*, v.42, no.2, 83–88 (2008).
- [8] Campillo A., Delgado F., Kiyek K. Gorenstein property and symmetry for one-dimensional local Cohen–Macaulay rings. *Manuscripta Mathematica*, v.83, no.3–4, 405–423 (1994).
- [9] Delgado F., Gusein-Zade S.M. Poincaré series for several plane divisorial valuations. *Proc. Edinb. Math. Soc. (2)*, v.46, no.2, 501–509 (2003).
- [10] Eisenbud D., Neumann W. *Three-Dimensional Link Theory and Invariants of Plane Curve Singularities*. Ann. of Math. Stud. 110, Princeton Univ. Press, Princeton, 1985.
- [11] Lipman J. Rational singularities with applications to algebraic surfaces and unique factorization. *Inst. Hautes Etudes Sci. Publ. Math.*, no.36, 195–279 (1969).
- [12] Némethi A. Poincaré series associated with surface singularities. *ArXiv: math.AG/0710.0987*.
- [13] Neumann W.D., Wahl J. Universal abelian covers of surface singularities. In: *Trends in singularities*, 181–190, Trends Math., Birkhäuser, Basel (2002).
- [14] Okuma T. Universal abelian covers of rational surface singularities. *J. London Math. Soc. (2)*, v.70, no.2, 307–324 (2004).
- [15] van Proeyen L., Veys W. The monodromy conjecture for zeta functions associated to ideals in dimension two. Preprint, Leuven University, 17 pp (2007).
- [16] Sabbah C. Modules d’Alexander et D -modules. *Duke Math. J.*, v.60, no.3, 729–814 (1990).
- [17] Zariski O., Samuel P. *Commutative Algebra, Vol 2*. van Nostrand, Princeton (1960).